

## **AMOXIMIN**

### **Injectable Suspension**

**Composition :**

Each 1 ml contains :

Amoxycillin Trihydrate 150 mg

**Description :**

Amoxycillin is a semi-synthetic penicillin with a broad-spectrum bactericidal activity against a large number of gram-positive and gram-negative micro-organisms .

The bactericidal action is due to inhibition of cell wall synthesis. Amoxycillin is mainly excreted in urine. A major part can also be excreted in bile.

**Indications :**

Gastrointestinal, respiratory and uro-genital infections caused by Amoxycillin sensitive micro-organisms.

Secondary bacterial infection in cases of viral diseases .  
(in cattle, goats, and sheep)

**Contraindication :**

- Hypersensitivity Amoxycillin.
- Animals with renal or liver failure.

**Side effects :**

Hypersensitivity reactions.

**Interactions :**

Should not be administered with tetracyclines, chloramphenicol, macrolides and lincosamides .

**Usage :**

For intramuscular injection

Shake well before use

**Dosage :**

The general dosage is: 1 ml per 15 kg body weight once a day for 3 days.

In sever cases the dose can be duplicated without fear of toxication .

**Withdrawal period :**

Meat : 14 days Milk : 3 days

**Storage conditions :**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

**Packing :**

Vial of 100 ml.