

## **CIPROMIN**

### **Oral Liquid**

**Composition:** each 1 ml contains:

Ciprofloxacin 100 mg

**Description:**

Ciprofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram negative bacteria like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp.

**Indications:**

Gastrointestinal, respiratory and urinary tract infections caused by Ciprofloxacin sensitive micro-organisms, like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella in ruminants and poultry .

**Contraindications:**

- Hypersensitivity to Ciprofloxacin
- Animals with hepatic or renal failure

**Side effects:**

Hypersensitivity reactions.

Administration to young animals can lead to arthropathy.

**Interactions:**

Avoid administration with tetracyclines, chloramphenicol, macrolides and lincosamides .

**Usage:**

Oral With Drinking Water

**Dosage:**

Poultry: 100-200 ml for 200 liter of drinking water for 3-4 days

Ruminants : 1 ml per 20 kg of body weight for 3-5 days

**Withdrawal period:**

Ruminants: Meat: 7 days      Milk: 3 days

Poultry: Meat: 4 days      Eggs: 3 days

**Storage conditions:**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

**Packing :** Bottle of 1 Liter.