

## **CLOMOJECT**

### **Sterile Powder For Injection**

#### **Composition :**

each Vial of 50 ml contains :

Amoxycillin Triglydrate    6 g

Cloxacillin Sodium         4 g

#### **Description :**

Cloxacillin is a specific antibiotic against micro-organisms which show resistance to Pencillin G like Staphylococcus which causes most Mastitis cases .

Amoxycillin is a semi-synthetic penicillin with a broad-spectrum bactericidal activity against a large number of gram-positive and gram-negative micro-organisms .

The bactericidal action is due to inhibition of cell wall synthesis. Amoxycillin is mainly excreted in urine. A major part can also be excreted in bile.

#### **Indications :**

Mastitis, Metritis, Gastrointestinal and uro-genital infections caused by Amoxycillin and/or Cloxacillin sensitive micro-organisms.

(in camels, horses, cattle, goats, and sheep)

#### **Contraindication :**

- Hypersensitivity Amoxycillin and / or Cloxacillin
- Animals with renal or liver failure.
- 

#### **Side effects :**

Hypersensitivity reactions.

#### **Usage :**

For intramuscular and intramammary injection

- Dilute the content of vial with sterilized water till 50 ml
- for intramammary injection : Milk out the udder completely, wash thoroughly with warm water containing suitable disinfectant.
- Shake well before use

#### **Dosage :**

##### Intramuscular injection :

Camels, horses and cattle : 25 ml per animal twice daily for 4 days

Sheep and goats : 7 ml per animal twice daily for 4 days

##### Intramammary injection :

Camels, horses and cattle: 10 ml per quarter or half for 3 days

Sheep and goats: 3-5 ml for per half for 3 days

#### **Withdrawal period :**

Meat : 30 days    Milk : 10 days

#### **Storage conditions :**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

#### **Packing :**

Vial of 50 ml.