

COXAZOL

Oral Liquid

Composition: each 1 ml contains

Sulphaquinoxaline sodium 100 mg

Diaveridine Hcl 20 mg

Description:

Sulfaquinoxaline is a chemotherapeutics with a bacteriostatic activity against many grampositive and gramnegative bacteria and has also coccidiostatic activity against Eimeria spp. It affects the metabolism of Folic Acid which is basic in the synthesis of the DNA of the Parasite cell (Eimeria). Diaveridine eases the toxicity of the sulfaquinoxaline , so we have an effective coccidiostat compound without side effects .

Indications:

- treatment and control of coccidiosis caused by Eimeria spp (tenella, acervulina, mevati, maxima, brunetti) in chicken, turkeys and rabbits .
- treatment of fowl cholera and fowl typhoid in chicken & turkeys.

Usage:

Oral With Drinking Water

Dosage:

1 ml per 1 liter for 3 days, stop for 2 days and resume for another 2 days

Precautions:

- not to be administrated for more than consecutive 3 days
- not to be used with Vitamin B complex

Withdrawal period:

Meat: 5 days, eggs: 3days.

Storage conditions:

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

Packing : Bottle of 1 Liter