

## ENROMIN 10%

### Oral Liquid

**Composition:** each 1 ml contains

Enrofloxacin 100 mg

**Description:**

Enrofloxacin belongs to the group of fluoroquinolones and acts bactericidal against both gram-positive bacteria and gram-negative bacteria. It is also active against Mycoplasmas.

**Indications:**

Gastrointestinal, respiratory and urinary tract infections, caused by micro-organisms susceptible to enrofloxacin (pasteurella, salmonella, mycoplasma, E-coli, Campylobacter, and Haemophilus) in Ruminants and Poultry.

**Contraindications:**

hypersensitivity to Enrofloxacin  
animals with hepatic or renal failure

**Side effects:**

Hypersensitivity reactions.

**Interactions:**

Not to be administered together with Tetracyclines, Chloramphenicol, Macrolides and Lincosamides.

**Usage:**

Oral With Drinking Water

**Dosage:**

Poultry: 100 ml per 200 liter for 3-4 days .

Ruminants: 1 ml per 20 kg body weight for 3-4 days.

**Withdrawal period :**

Ruminants: Meat: 7 days Milk: 3 Days

Poultry: Meat: 4 days Eggs: 3 days.

**Storage conditions:**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight

**Packing:** Bottle of 1 Liter and 100 ml .