

LINCOJECT

Injectable Solution

Composition :

Each ml contains:

Spectinomycin Sulphate	100 mg
Lincomycin Hydrochloride	50 mg

Description :

The combination of lincomycin and spectinomycin acts additive and in some cases synergistic. Spectinomycin acts bacteriostatic or bactericidal, depending on the dose, against mainly Gram-negative bacteria like Campylobacter, E. coli, Mycoplasma and Salmonella . Lincomycin acts bacteriostatic against mainly Gram-positive bacteria like Mycoplasma, Staphylococcus, Streptococcus .

Indications :

Gastrointestinal and respiratory infections caused by lincomycin and spectinomycin sensitive micro-organisms, like Campylobacter, E. coli, Mycoplasma, Salmonella, Staphylococcus, and Streptococcus, in horses, camels, cattle, cats, dogs, goats, sheep, poultry and turkeys

Contraindications:

- Hypersensitivity to Spectinomycin and/or Lincomycin .
- Animals with renal or liver failure .
- Not to be used for hens producing eggs for human consumption.

Side effects :

Hypersensitivity reaction .

Interactions:

Oxytetracycline should not be used concomitantly with Penicillins, Cephalosporines or Quinolones .

Usage :

Poultry and Turkey : subcutaneous injection

Other animals : intramuscular injection

Dosage :

Large animals : 1 ml per 10 kg of body weight, for 3 - 5 days.

Small animals : 1 ml per 5 kg of body weight, for 3 - 5 days.

Withdrawal Period :

Poultry and turkey : Meat : 7 days .

Other animals : Meat : 14 days Milk : 3 days .

Storage conditions :

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

Packing :

Vial of 100 ml .