

LINCOMIN
Water Soluble Powder

Composition :

Each jar of 100 g contains:

Spectinomycin Sulphate	44.5 g
Lincomycin Hydrochloride	22.2 g

Description :

The combination of lincomycin and spectinomycin acts additive and in some cases synergistic. Spectinomycin acts bacteriostatic or bactericidal, depending on the dose, against mainly Gram-negative bacteria like Campylobacter, E. coli, Mycoplasma and Salmonella . Lincomycin acts bacteriostatic against mainly Gram-positive bacteria like Mycoplasma, Staphylococcus, Streptococcus .

Indications :

In Poultry for the treatment and prevention of Chronic respiratory disease CRD and to increases the rate of body gain.

Contraindications:

- hypersensitivity to Spectinomycin and/or Lincomycin .
- animals with renal or liver failure .
- not for hens producing eggs for human consumption.

Side effects :

Hypersensitivity reaction .

Interactions:

Should not be used concomitantly with Penicillins, Cephalosporines or Quinolones .

Usage :

Oral with drinking water

Dosage :

100 g per 125 liter of drinking water, give as the sole source of drinking water for flocks of 5-7 days old .

Withdrawal Period :

Meat : 7 days .

Storage conditions :

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

Packing :

Jar of 100 g .