## **Water Soluble Powder**

## **Composition:**

Each sachet of 100 g contains:

Sulphadimidine Sodium 12.5 g Sulphathiazole Sodium 12.5 g Sulphadiazole Sodium 12.5 g

## **Description:**

All sulphanomides acts against the synthesis of Para Amino Benzoic Acid which is essential in the synthesis of the DNA of the microbe cell .

Sulphonamides also acts against Folic Acid which is essential in the microbe metabolism.

The combination of these three sulphanomides results in a wide range of effect against broad spectrum of gram- positive and gram- negative microorganisms.

## Indications:

Gastrointestinal, respiratory and urogenital infections caused by sulphanomides sensitive microorganisms like E.coli, Staphylococcus, Streptococcus, Pasteurella, Salmonela in camel, cattle, sheep, goats, and poultry

## **Contraindications:**

- hypersensitivity to sulfonamides.
- animals with a renal or Hepatic failure.

### Usage:

Oral with drinking water

#### Dosage:

Camels, cattle, sheep and goats: 1g per 10 kg of body weight.

Poultry: 200 g per 200 liter of drinking Water.

### Precaution:

Not to be administrated for more than 5 days because of the accumulation of the sulphanomides in the kidney .

### Withdrawal Period:

<u>Ruminants:</u> Meat : 5 days Milk : 3 days <u>Poultry:</u> Meat :5 days eggs : 4 days

# **Storage conditions:**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

### Packing:

Sachet of 100 g.