

STREPTOMIN

Injectable Suspension

Composition :

Each 1 ml contains :

Procaine Penicillin G 200 000 IU

Dihydrostreptomycin Sulphate 200 mg

Description :

The combination of procaine penicillin G and Dihydrostreptomycin acts additive and in some cases synergistic. Procaine Penicillin G is a small-spectrum penicillin with a bactericidal action against mainly gram-positive bacteria .

Dihydrostreptomycin is an aminoglycoside with a bactericidal action against mainly gram-negative bacteria.

Indications :

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms., in cattle, goats, and sheep .

Contraindication :

- Hypersensitivity to penicillin, procaine and aminoglycosides.
- Animals with renal or liver failure.

Side effects :

Hypersensitivity reactions.

Interactions :

Should not be administered with tetracyclines, chloramphenicol, macrolides and lincosamides .

Usage :

For intramuscular injection

Shake well before use

Dosage :

Cattle : 1 ml per 20 kg body weight for 3 days.

Calves, sheep and goats : 1 ml per 10 kg body weight for 3 days

Withdrawal period :

Meat : 21 days Milk : 3 days

Storage conditions :

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

Packing :

Vial of 100 ml.