

SULPHAMIN 10%

Oral Suspension

Composition: each ml contains:

Sulphadiazine Sodium	100 mg
Trimethoprim	20 mg

Description:

The combination of Trimethoprim and Sulphadiazine acts synergistically against a large number of gram-positive and gram-negative bacteria . After oral administration, there will be a rapid absorption of both components from the gastrointestinal tract.

Indications:

Gastrointestinal and respiratory infections caused by trimethoprim and sulfadiazine sensitive micro-organisms, like E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus., in camel, cattle, sheep, goat and poultry .

Contraindication:

- Hypersensitivity to trimethoprim and/or sulphonamides.
- Animals with a renal and/or hepatic failure .

Side effects:

- Hypersensitivity reaction .
- After long-term treatment and high dosages crystalluria can occur.

Usage:

Oral With Drinking Water

Dosage:

Ruminants: 2 ml per 10 kg Body Weight for 4–5 days

Poultry: 200 ml per 100 liters of drinking water for 4-5 days.

Withdrawal Period:

Poultry: Meat:3 days Eggs: 3 days .

Ruminant: Meat: 8 days Milk: 3 days

Storage conditions: Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

Packing: Bottle of 1 liter – 500 ml – 200 ml and 100 ml .