

TYLOPRIM PLUS

Oral Suspension

Composition: each ml contains:

Sulphadiazine Sodium	175 mg
Trimethoprim	35 mg
Tylosin Tartrate	63 mg

Description:

Tylosin, a macrolide antibiotic, has its bactericidal action by inhibiting the protein synthesis of the bacteria cell

It is characterized by its rapid absorption from the digestive tract, and after oral administration, therapeutically active blood concentrations of Tylosin are reached within 15- 30 minutes and then to other tissues (Lungs, air sacs, ovary and urinary tract).

is active against particularly gram-positive bacteria, mycoplasmas, and Chlamydia

The combination of Trimethoprim and Sulphadiazine acts synergistically against a large number of gram-positive and gram-negative bacteria . After oral administration, there will be a rapid absorption of both components from the gastrointestinal tract.

This medicine contains above three active components and affects a broad spectrum of Positive and negative bacteria besides Mycoplasma and Chlamydia.

Indications:

TYLOPRIM PLUS is used in control and treatment of gastrointestinal and respiratory infections in poultry especially CRD, Coryza and Avian Cholera

Contraindications:

Avoid administration to poultry producing eggs for human consumption.

Usage:

Shake well before use
Oral With Drinking Water

Dosage:

Poultry: 1-2 ml per 1 liters of drinking water for 4-8 days.

Withdrawal Period:

Poultry: Meat: 10 days

Storage conditions:

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

Packing:

Bottle of 500 ml and 100 ml .