

## **TYMOJECT**

### **Injectable Solution**

**Composition :**

Each ml contains:

Tylosin Tartrate            200 mg

**Description :**

Tylosin, a macrolide antibiotic, is active against particularly gram-positive bacteria, some Spirochetes (including Leptospira), Actinomyces, Mycoplasmas, Haemophilus pertussis, and some gram-negative cocci. After parenteral administration, therapeutically active blood concentrations of Tylosin are reached within two hours.

**Indications :**

Infections caused by micro-organisms susceptible to Tylosin : (in cattle, calves, sheep, goats)

- respiratory tract infections.
- dysentery
- arthritis
- mastitis
- endometritis.
- shipping fever
- pneumonia
- foot rot
- bronchitis
- post surgery operation infections

**Contraindications:**

Hypersensitivity to Tylosin .

**Interactions:**

Avoid administration with penicillines, cephalosporines, or quinolones

**Usage :**

For intramuscular injection

**Dosage :**

The general dose is: 1 ml per 20 kg body weight daily for 3-5 days.

**Withdrawal Period :**

Meat : 10 days    Milk : 4 days .

**Storage conditions :**

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight .

**Packing :**

Vial of 100 ml .