

ZAMIN

Oral Suspension

Composition: each 1 ml contains

Oxyclozanide 60 mg

Levamisole Hydrochloride 30 mg

Description:

Levamisole and oxyclozanide act against a broad spectrum of gastrointestinal worms and against lung worms.

Levamisole is a broad-spectrum anthelmintic and active against adult and most larval stages of intestinal nematodes and lungworms in Ruminants.

Oxyclozanide is a salicylanilide and acts against Trematodes, and bloodsucking nematodes .

Indications:

Prevention and treatment of:

Gastrointestinal nematodes: Trichostrongylus, Cooperia, Ostertagia, Haemonchus, Nematodirus, Chabertia, Bunostomum,

lungwormin: Dictyocaulus

liverfluke: Fasciola Hepatica .

Contraindications:

- Avoid administration of during early pregnancy.
- Animals with renal or hepatic failure .
- Hypersensitivity to oxyclozanide and/or levamisole .

Side effects:

Overdosages can cause excitation, lachrymation, sweating, excessive salivation, coughing, hyperpnoea, vomiting, colic and spasms.

Usage:

For oral administration

Shake well before use

Dosage:

Cattle, calves: 2.5 ml per 10 kg of body weight.

Sheep and goats: 1 ml per 4 kg of body weight.

Withdrawal period:

Meat: 28 days, Milk: 4 days.

Storage conditions:

Store in airtight containers, in a moderate temperature (15-25 C) and protect from direct sunlight.

Packing: Bottle of 1 Liter - 500 ml and 100 ml .